

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 June 2002 (27.06.2002)

PCT

(10) International Publication Number
WO 02/50017 A1

(51) International Patent Classification⁷: **C07C 213/02,**
217/74

(FR). **STUDER, Martin** [CH/CH]; Im Zimmerhof 2,
CH-4054 Basel (CH). **BURKHARDT, Stefan** [CH/CH];
Gansacherweg 42, CH-4460 Gelterkinden (CH).

(21) International Application Number: **PCT/EP01/14604**

(22) International Filing Date:
12 December 2001 (12.12.2001)

(74) Common Representative: **CIBA SPECIALTY CHEMI-**
CALS HOLDING INC.; Patentabteilung, Klybeckstrasse
141, CH-4057 Basel (CH).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1101/MAS/00 20 December 2000 (20.12.2000) IN
390/MAS/01 14 May 2001 (14.05.2001) IN

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **CIBA**
SPECIALTY CHEMICALS HOLDING INC. [CH/CH];
Klybeckstrasse 141, CH-4057 Basel (CH).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **PAINGANKAR,**
Niranjan [IN/IN]; Dhanvantary Niwas, 1st Floor, Subbhash
Road, Vile Parle East, Mumbai 400 057 (IN). **MUM-**
BAIKAR, Vilas, N. [IN/IN]; 202 Vindhychal CHS, Lok
Everest complex, ACC Road, Mulund West, Mumbai
400 081 (IN). **EKKUNDI, Vadiraj, S.** [IN/IN]; D-318
Ciba Housing Colony, Goregaon East, Mumbai 400
063 (IN). **JALETT, Hans-Peter** [CH/CH]; Solothurn-
erstrasse 2, CH-4143 Domach (CH). **SIEGRIST, Urs**
[CH/CH]; Rebenweg 32, CH-5074 Eiken (CH). **VAN**
DER SCHAAF, Paul, Adriaan [NL/CH]; Marsstrasse
17, CH-4123 Allschwil (CH). **BIENEWALD, Frank**
[DE/FR]; 63, rue de Hagenthal, F-68220 Hegenheim

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: **PROCESS FOR THE PREPARATION OF PHENETHYLAMINE DERIVATIVES**

(57) Abstract: Disclosed is a process for the preparation of a compound of formula wherein R₁ is hydrogen, hydroxyl, or unsubstituted or substituted alkyl or alkoxy, R₂ is hydrogen or a substituent which can be converted to hydrogen, and n is 0, 1 or 2, comprising hydrogenating a compound of formula wherein R₁, R₂ and n are as defined above, in the presence of a nickel or cobalt catalyst.

WO 02/50017 A1

Process for the preparation of phenethylamine derivatives

The present invention is directed to a process for the preparation of phenethylamine derivatives by hydrogenation of phenylacetonitriles in presence of a nickel or cobalt catalyst.

The compounds of formula (1) are known for being particularly useful as synthesis intermediates for preparing pharmaceutical active substances which are central nervous system antidepressants. An important substance is Venlafaxine (see Merck Index Twelfth Edition 1996, No. 10079). The preparation of this compound is described in US-A-4,535,186.

According to US-A-4,535,186, Example 2, intermediates of formula (1) are prepared by hydrogenation in the presence of a rhodium catalyst. The use of rhodium catalysts provides economical drawbacks and therefore such catalysts would have to be recycled, resulting in a further process step and the possibility of varying catalytic effect of the recycled catalyst.

It is the object of the present invention to provide a process for the preparation of phenethylamine derivatives in high yields which meets economical demands.

The present invention relates to a process for the preparation of a compound of formula




or salts thereof, wherein

R₁ is hydrogen, hydroxyl, or unsubstituted or substituted alkyl or alkoxy,

R₂ is hydrogen or a substituent which can be converted to hydrogen, and

n is 0, 1 or 2,

comprising hydrogenating a compound of formula



(2),

wherein R_1 , R_2 and n are as defined above, in the presence of a nickel or cobalt catalyst.

R_1 as alkyl can be a straight-chain or branched alkyl substituent, preferably C_1 - C_4 alkyl, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, *tert*-butyl. The substituents mentioned can be unsubstituted or substituted. An example for such substituents is phenyl.

R_1 as alkoxy can be a straight-chain or branched alkoxy substituent, preferably C_1 - C_4 alkoxy, such as methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-butoxy, *sec*-butoxy, *tert*-butoxy. Methoxy is preferred. The substituents mentioned can be unsubstituted or substituted. An example for such substituents is phenyl.

Preferably, R_1 is hydroxyl or C_1 - C_4 alkoxy, especially C_1 - C_4 alkoxy. Methoxy is highly preferred. It is preferred that R_1 is bonded to the benzene ring in *para*-position.

Examples for substituents R_2 which can be converted to hydrogen are silyl, benzyl, formyl or C_2 - C_6 alkanoyl. The conversion can be carried out according to known methods.

Preferably, R_2 is hydrogen, formyl or C_2 - C_6 alkanoyl, especially hydrogen.

For n the number 1 is preferred.

According to a preferred embodiment of the present invention R_1 is methoxy, R_2 is hydrogen and n is 1.

According to the present invention nickel or cobalt catalysts well known and used in the hydrogenation field can be used.

For example, a nickel catalyst can be made by thermally decomposing nickel formate or other heat-labile nickel salts, for example in fatty oil, or by precipitating a nickel salt on an

inert carrier followed by a reduction with hydrazine or hydrogen gas. A nickel catalyst also can be prepared by the treatment of electrolytically precipitated nickel hydroxide which may be prepared by passing direct current through a cell using nickel as the anode and using the dilute solution of an alkali salt to the weak acid as an electrolyte. The nickel hydroxide so prepared may be conventionally reduced, such as, in the presence of hydrogen gas or hydrazine.

The nickel or cobalt catalyst also may be promoted as is conventional in this field. So, the catalysts may contain, for example, Group VIB metals, Group VIB metal compounds which are reducible by hydrogen to the corresponding elemental metal, or manganese or iron promoters. Specific examples of the group VIB metals or hydrogen-reducible metal compounds include elemental chromium, chromium acetate, chromium chloride, chromium oxide, elemental molybdenum, molybdenum hydroxide, molybdenum oxide, elemental tungsten, tungsten chloride, tungsten oxide, and the like, and mixtures of any two or more thereof. The weight ratio of the group VIB component to the metal component of the nickel or cobalt catalyst can be any suitable value, but will generally be in the range from about 0.001:1 to about 0.2:1 and preferably in the range from about 0.005:1 to about 0.1:1.

The catalysts can be in supported or unsupported form. Typical support materials include, for example, carbon, aluminium oxide, silicium dioxide, Cr_2O_3 , titanium dioxide, zirconium dioxide, zinc oxide, calcium oxide, magnesium oxide, barium sulfate, calcium carbonate or aluminium phosphate. The nickel or cobalt catalyst can be bound on the substrate in an amount of, for example, about 1.0 - 20.0 % by weight.

The preferred catalysts are Raney nickel and Raney cobalt catalysts. Such catalysts are, for example, formed by mixing nickel and aluminum or cobalt and aluminum and subsequently treating the respective mixtures with a suitable base, such as sodium hydroxide to remove the aluminum, thus leaving a highly reactive nickel or cobalt metal catalyst.

In all cases nickel catalysts are preferred; highly preferred are Raney nickel catalysts.

According to a preferred embodiment of the present invention the nickel or cobalt catalysts are pretreated with a carboxylic acid or a salt or an anhydride thereof or an ammonium salt,

or a vanadium-, a tungsten-, or a molybdenum compound. Mixtures of at least two of these compounds can also be used.

Carboxylic acids, including polycarboxylic acids, having from 1 to 8 carbon atoms, especially 2 to 8 carbon atoms, are preferred. The carboxylic acids can be unsubstituted or substituted by, for example, hydroxy or halogen (like fluorine). Examples of such carboxylic acids are formic acid, acetic acid, propionic acid, butanoic acid, pentanoic acid, oxalic acid, malonic acid, succinic acid, maleic acid, malic acid, citric acid, tartaric acid, trifluoroacetic acid. Examples for anhydrides are acetic anhydride, propionic anhydride and further anhydrides of the carboxylic acids mentioned above. Examples for salts of the carboxylic acids are the corresponding alkali metal salts, like sodium salts.

Preferred are carboxylic acids having from 2 to 4 carbon atoms, especially 2 carbon atoms. These carboxylic acids can be unsubstituted or substituted as given above. This definition should also include the salts and anhydrides thereof. Highly preferred is acetic acid (as well as the salts and the anhydride thereof).

Ammonium salts are, for example, ammonium halides, like ammonium chloride or especially ammonium fluoride

Preferred vanadium compounds are those in which vanadium has the oxidation state 0, 2, 3, 4 or 5. Elemental vanadium is also suited. Examples of such compounds are V_2O_5 , $VOCl_3$, V_2O_4 , NH_4VO_3 . Specially preferred are acetyl acetonate (acac) containing compounds, examples of such compounds are $V(acac)_3$, $VO(acac)_2$. The vanadium compounds are used in catalytic amounts, solved or dispersed.

Preferred tungsten or molybdenum compounds are compounds in which the metal has the oxidation state 0, 2, 3, 4, 5 or 6. Examples are H_2WO_4 , $H_3[P(W_3O_{10})_4]$, H_2MoO_4 , $H_3[P(Mo_3O_{10})_4]$.

For the pretreatment of the nickel or cobalt catalysts the use of carboxylic acids or salts or anhydrides thereof, or acetyl acetonate containing compounds is preferred. Highly preferred are carboxylic acids.

Advantageously the pretreatment is carried out in an aqueous medium. As a rule the aqueous medium contains the nickel or cobalt catalyst and the agent used for the treatment. Usually the treatment is carried out at ambient temperature, although also lower or higher temperatures can be used.

The reaction conditions of the hydrogenation, like time, temperature and pressure, are to a large extent interchangeable, as will be recognized by the skilled practitioner.

Preferably, the hydrogenation is carried out in an organic solvent, like an alcohol (for example methanol or especially ethanol). Furthermore, it is preferred that the hydrogenation is carried out in presence of a base (examples are NH_3 , NH_4OH and NaOH).

The cited catalysts can be present in the reaction mixture in an amount of about 0.1 to 500% by weight, preferably of 20 to 200% by weight, based on the amount of educt used.

The educts can be present in the reaction mixture in an amount of about 0.1 to 80% by weight, preferably of 5 to 20% by weight, based on the weight of the reaction mixture.

The temperature to be used for the reaction can be, for example, 0 to 200°C, a temperature of 20 to 120°C being preferred. Highly preferred is a temperature of 20 to 80°C.


The reaction time can vary, for example, from 0.1 to 24 hours. Usually the reaction time is from 0.1 to 14 hours, especially from 0.1 to 4 hours.

The hydrogen pressure is, for example, 1 to 200 bar, especially 1 to 100 bar. A hydrogen pressure of 20 to 60 bar is preferred.

The catalyst can be re-used 1 to 100 times, especially 1 to 10 times; surprisingly, it is possible to re-use the catalyst without further treatment and the recycled catalyst shows no decrease in activity or selectivity.

After the reaction the product can be further purified according to known methods. For example, the product obtained after the reaction can be dissolved in an organic solvent (like diisopropyl ether) and separated by addition of an acid and filtration of the corresponding


salt. Preferably, the product can be purified by addition of a hydrogen halide (like hydrogen chloride) or a carboxylic acid having from 1 to 6 carbon atoms (like formic acid). Highly preferred is the addition of formic acid after which the compound is separated by filtration as the compound of formula (1a)



A further object of the present invention are compounds of formula (1a). As to these compounds the meanings and preferences given above apply. The compounds of formula (1a) do not require special treatment when they are further converted into compounds of formula (3).

Alternatively, the product can be used directly, without purification, for further reactions like those described below.

Another object of the present invention is a process for the preparation of a compound of formula



wherein

R_1 , R_2 and n are as defined above,

comprising hydrogenating a compound of formula



in the presence of a nickel or cobalt catalyst to give the compound of formula




and converting the compound of formula (1) to the compound of formula (3).

As to R_1 , R_2 and n the above meanings and preferences apply.

The conversion of the compound of formula (1) to the compound of formula (3) can be carried out according to known processes. Such a conversion and the reaction conditions to be used are described in US-A-4,535,186 (see especially Example 3).

In general, a method for such a conversion comprises the following steps:



and B) preparation of the corresponding hydrochloride salts.

Step A) can, for example, be carried out by reaction of the compound of formula (1) with formaldehyde, formic acid in a large excess of water. Step B) can be carried out in

conventional manner by forming the acid addition salt, whereby preferably an equimolar amount of hydrochloric is used.

According to the present invention the intermediates of formula (1) can be obtained in high yields. The use of expensive catalysts can be dispensed with.

The following examples illustrate the invention:

Example 1:

a) Pretreatment of the catalyst

135 g of Raney nickel are washed three times each with 135 ml of water followed by 250 ml of an aqueous solution of acetic acid (5% by volume) and again three times each with 1000 ml of water.

b) Hydrogenation

A 5 liter autoclave is charged with 180 g of 1-[cyano(4-methoxyphenyl)methyl]cyclohexanol, 2400 ml of methanol, 600 ml of aqueous ammonia (25% by volume) and 135 g of Raney nickel (pretreated as given under 1a)) and the mixture is subjected to hydrogenation at 27 to 30°C and 120 psi pressure of hydrogen for 9 to 10 hours. The reaction mixture is filtered through 100 g of the celite and the catalyst bed is washed with 700 ml of methanol. The filtrate is concentrated to get 167.1 g of the crude product as an oily residue. 1.1 g of the crude product is dissolved in 2 ml of dry ethyl acetate and about 2 ml of isopropanolic HCl are added (the pH of the solution is about 2). The solvent is removed at high vacuum to give 1.14 g of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol hydrochloride.

Example 2:

a) Pretreatment of the catalyst

50 ml of an aqueous solution of acetic acid (5% by volume) are added to 5 g of Raney nickel (suspension containing 60% by weight of the catalyst). The resulting suspension is stirred at ambient temperature for 10 to 15 minutes. After separation the catalyst is washed four times with deionised water.

b) Hydrogenation

An autoclave is charged with 1.2 g of 1-[cyano(4-methoxyphenyl)methyl]cyclohexanol, the amount of Raney nickel obtained after the pretreatment given above under 2a), 24 ml of ethanol and 6 ml of aqueous ammonia (25% by volume) and the mixture is subjected to hydrogenation at 60°C and 40 bar pressure of hydrogen for 140 minutes. The catalyst is filtered off and washed with ethanol. The filtrate and the ethanol used to wash the catalyst are combined and concentrated by evaporation to get 1.2 g of the crude product as an oily residue. The crude product contains 79% by weight of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol (HPLC). The crude product obtained as given above is dissolved in diisopropyl ether and HCl is added in order to form the HCl addition salt. The salt obtained is filtered off, washed with diisopropyl ether and dried in vacuum. 1.23 g of white crystals are obtained having a melting point of 169°C and containing 84% by weight of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol hydrochloride (HPLC).

Example 3:

Example 2 is repeated, using twice the amount of pretreated Raney nickel given in Example 2, carrying out the hydrogenation for 70 minutes instead of 140 minutes and dispensing with the formation of the HCl addition salt. 1.2 g of the crude product are obtained containing 86% by weight of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol (HPLC).

Example 4:

Example 2 is repeated, using the same amount of Raney cobalt instead of Raney nickel, carrying out the hydrogenation for 90 minutes instead of 140 minutes and dispensing with the formation of the HCl addition salt. 1.2 g of the crude product are obtained containing 92% by weight of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol (HPLC).

Example 5:**a) Pretreatment of the catalyst**

A saturated aqueous solution of V(acac)₃ [acac = acetyl acetonate] in 500 ml of deionised water is added to 50 g of Raney nickel (suspension containing 60% by weight of the catalyst). The resulting suspension is stirred at ambient temperature for 15 to 20 minutes. After separation the catalyst is washed five times with 500 ml of deionised water.

b) Hydrogenation

An autoclave is charged with 33 g of 1-[cyano(4-methoxyphenyl)methyl]cyclohexanol, the amount of Raney nickel catalyst obtained after the pretreatment given above under 5a), 320 ml of ethanol and 80 ml of aqueous ammonia (25% by volume) and the mixture is subjected to hydrogenation at 60°C and 40 bar pressure of hydrogen for 640 minutes. The catalyst is filtered off and washed with ethanol. The filtrate and the ethanol used to wash the catalyst are combined and concentrated by evaporation to get 33 g of the crude product as an oily residue. The crude product contains 93% by weight of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol (HPLC).

The crude product can be processed directly to give the corresponding HCl addition salt.

Example 6:**a) Pretreatment of the catalyst**

50 ml of an aqueous solution of $H_3[P(W_3O_{10})_4] \cdot xH_2O$ (0.5% by volume) is added to 5 g of Raney nickel (suspension containing 60% by weight of the catalyst). The resulting suspension is stirred at ambient temperature for 15 to 20 minutes. After separation the catalyst is washed five times with deionised water.

b) Hydrogenation

An autoclave is charged with 1.2 g of 1-[cyano(4-methoxyphenyl)methyl]cyclohexanol, the amount of Raney nickel obtained after the pretreatment given above under 6a), 24 ml of ethanol and 6 ml of aqueous ammonia (25% by volume) and the mixture is subjected to hydrogenation at 60°C and 40 bar pressure of hydrogen for 300 minutes. The catalyst is filtered off and washed with ethanol. The filtrate and the ethanol used to wash the catalyst are combined and concentrated by evaporation to get 1.2 g of the crude product as an oily residue. The crude product contains 90% by weight of 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol (HPLC).

Example 7:**a) Formation of the 1-[2-amino-1-(methoxyphenyl)ethyl]cyclohexanol formic acid salt:**

49.8 g of crude 1-[2-amino-1-(methoxyphenyl)ethyl]cyclohexanol from the nitrile hydrogenation was suspended in 250 ml ethylacetate and 12 ml formic acid (98-100 %) were added in one portion. The mixture was heated to reflux under stirring and subsequently

cooled down to room temperature. The product was filtered off, washed with 100 ml hexane and dried in vacuo, yielding 48.1 g (82 %) of crude formiate salt.

b) Purification of the formiate salt:

The crude formiate salt was suspended in 1 l ethylacetate, heated to reflux for 1 hour, cooled to room temperature and filtered off. At this stage, the product has got already a much improved purity as shown by HPLC (> 98 %). The purification procedure was repeated one time and the final product was dried in vacuo at 40°C, giving 29.6 g product (50 %) as white crystalline. The purity was determined by HPLC to be > 98 %.

Example 8:

Reductive methylation of 1-[2-amino-1-(methoxyphenyl)ethyl]cyclohexanol formic acid salt: 29.6 g of the formiate salt from above was mixed with 100 ml water, 17 ml formic acid (98-100 %) and 26 ml formaldehyde (37 %) and heated to reflux for 20 hours. The mixture was cooled to room temperature, acidified with 25 ml 4 N hydrochloride acid to pH < 1 and extracted five times with 50 ml ethylacetate. The pH of the aqueous phase was adjusted to > 12 with 30 % aqueous sodium hydroxide and 150 ml toluene were added. The mixture was passed through a R3 filter, the phases were separated and the aqueous phase was again extracted with 50 ml toluene. The combined organic phases were washed with 100 ml water. Under stirring 30 ml 4.2 N hydrochloride acid in 1,4-dioxane were added over 5 minutes and the formed suspension was stirred for additional 1 hour. The product was filtered off, washed two times with 50 ml hexane and dried in vacuo at 40°C yielding 26.9 g of the hydrochloride salt as white crystalline (86 %), pure by HPLC.

Example 9:

a) Pretreatment of the catalysts

6.00 kg Raney Nickel are placed in a 50 l steel autoclave. 40 l of water containing 115.2 g VO(acac)₂ are added. The mixture is stirred for 30 minutes at room temperature. Then the aqueous phase is removed.

b) Hydrogenation

4.00 kg of 1-[cyano(4-methoxyphenyl)methyl]cyclohexanol is suspended in 5 l EtOH and transferred to the autoclave containing the modified Raney Nickel. Then, 22 l of EtOH are added and the autoclave is closed and checked for leaks. After this, the 6 l of 25% NH₄OH in

water are added through an addition port. Then the gases in the autoclave are displaced by nitrogen (3 times) and hydrogen (3 times). Then the autoclave is pressurized to 40 bar and the reaction is started by turning the stirrer on. The reaction mixture is heated to 60 °C in about 20 minutes and the reaction is continued for 2 hours at 60°C. Then the autoclave is cooled to room temperature, the hydrogen is displaced by nitrogen and the autoclave is opened. After filtration of the catalyst, the reaction mixture is evaporated to dryness. HPLC analysis shows a selectivity for the desired 1-[2-amino-1-(4-methoxyphenyl)ethyl]-cyclohexanol] of 93% (HPLC).

Example 10:

a) Pretreatment of the catalysts

22.5 g Raney Nickel are placed in a 300 ml steel autoclave equipped with a filtration frit. 220 ml of an aqueous solution saturated with V(acac)₃ is added and stirred during 30 minutes. After standing over night, the water is discharged through the frit and the catalysts is washed with 50 ml water and 3 x 50 ml EtOH.

b) Hydrogenation

15 g of 1-[cyano(4-methoxyphenyl)methyl]cyclohexanol] is suspended in 50 ml EtOH and transferred to the 0.3 l autoclave containing the modified Raney Nickel. Then, 50 ml of EtOH are added and the autoclave is closed and checked for leaks. After this, the 30 ml of 25% NH₄OH in water are added through an addition port. Then the gases in the autoclave are displaced by nitrogen (3 times) and hydrogen (3 times). Then the autoclave is pressurized to 42 bar and the reaction is started by turning the stirrer on. The reaction mixture is heated to 60 °C in about 20 minutes and the reaction is continued for 2 hours at 60°C. Then the autoclave is cooled to room temperature, the hydrogen is displaced by nitrogen and the autoclave is opened. After filtration of the catalyst, the reaction mixture is evaporated to dryness. HPLC analysis shows a selectivity for the desired 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol] of 86% (HPLC).

c) Hydrogenation: Catalyst re-use

The catalyst from the experiment described above is washed with 50 ml EtOH, then, the hydrogenation experiment is repeated exactly as described above. The same reaction times are observed. HPLC analysis shows a selectivity for the desired 1-[2-amino-1-(4-methoxyphenyl)ethyl]cyclohexanol] of 85%. The selectivity observed in subsequent runs is

- 13 -

between 85% and 87% (HPLC) and no noticeable catalyst deactivation is observed over four re-uses. The Vanadium content of the products is always < 2 ppm, and the Nickel content varies between 1 and 3 ppm.

Claims

1. A process for the preparation of a compound of formula



or salts thereof, wherein

R_1 is hydrogen, hydroxyl, or unsubstituted or substituted alkyl or alkoxy,

R_2 is hydrogen or a substituent which can be converted to hydrogen, and

n is 0, 1 or 2,


comprising hydrogenating a compound of formula



wherein R_1 , R_2 and n are as defined above, in the presence of a nickel or cobalt catalyst.

2. A process according to claim 1, wherein R_1 is hydroxy or C_1 - C_4 alkoxy which is unsubstituted or substituted by phenyl, preferably methoxy.
3. A process according to claim 1 or 2, wherein R_1 is bonded in para position.
4. A process according to any of claims 1 to 3, wherein R_2 is hydrogen, silyl, benzyl, formyl or C_2 - C_6 alkanoyl, preferably hydrogen.
5. A process according to any of claims 1 to 4, wherein n is 1.
6. A process according to claim 1, wherein R_1 is methoxy, R_2 is hydrogen and n is 1.


7. A process according to any of claims 1 to 6, wherein a nickel or cobalt catalyst is used which has been pretreated with a carboxylic acid or a salt or an anhydride thereof, or with an ammonium salt, or a vanadium compound or a tungsten- or a molybdenum compound, or a mixture of at least two of these compounds.
8. A process according to claim 7, wherein a carboxylic acid having from 1 to 8 carbon atoms is used, preferably acetic acid.
9. A process according to claim 7, wherein an acetyl acetate containing vanadium compound is used.
10. A process according to any of claims 7 to 9 wherein the pretreatment is carried out in an aqueous medium.
11. A process according to any of claims 1 to 10, wherein the catalyst is Raney nickel or Raney cobalt.
12. A process according to any of claims 1 to 11, wherein a nickel catalyst, preferably Raney nickel, is used.
13. A process according to any of claims 1 to 12, wherein the hydrogenation is carried out in presence of an organic solvent, preferably an alcohol.
14. A process according to any of claims 1 to 13, wherein a recycled catalyst is used.
15. A process according to claim 1, wherein after hydrogenation the product is purified by reaction with formic acid forming the compound of formula



(1a),


wherein R_1 , R_2 and n are as defined in claim 1.

16. A compound of the formula



wherein R_1 , R_2 and n are as defined in claim 1.

17. A process for the preparation of a compound of formula



wherein

R_1 is hydrogen, hydroxyl, or unsubstituted or substituted alkyl or alkoxy,


R_2 is hydrogen or a substituent which can be converted to hydrogen, and

n is 0, 1 or 2,

comprising hydrogenating a compound of formula



wherein R_1 , R_2 and n are as defined above, in the presence of a nickel or cobalt catalyst to give the compound of formula



(1),

wherein R_1 , R_2 and n are as defined above, and converting the compound of formula (1) to the compound of formula (3).

18. A process according to claim 17, wherein a nickel or cobalt catalyst is used which has been pretreated with a carboxylic acid or a salt or an anhydride thereof, or with an ammonium salt, or a vanadium compound or a tungsten- or a molybdenum compound, or a mixture of at least two of these compounds.

19. A process according to claim 17 or 18, wherein R_1 is methoxy, R_2 is hydrogen and n is 1.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 01/14604

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C213/02 C07C217/74

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 535 186 A (HUSBANDS G E MORRIS ET AL) 13 August 1985 (1985-08-13) cited in the application column 1 -column 3; claims; example 2 ---	1-19
A	YARDLEY J P ET AL: "2-PHENYL-2-(1-HYDROXYCYCLOALKYL)ETHYLAMINE DERIVATIVES: SYNTHESIS AND ANTIDEPRESSANT ACTIVITY" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 33, 1990, pages 2899-2905, XP000891765 ISSN: 0022-2623 page 2900 -page 2901 -----	1-19

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

24 April 2002

Date of mailing of the international search report

14/05/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Sánchez García, J.M.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

.../EP 01/14604

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4535186	A	13-08-1985	AT 28628 T	15-08-1987
			AU 567524 B2	26-11-1987
			BG 60659 B2	30-11-1995
			CA 1248540 A1	10-01-1989
			DE 3372753 D1	03-09-1987
			DK 571383 A ,B,	14-06-1984
			EG 17630 A	30-06-1992
			EP 0112669 A2	04-07-1984
			ES 527938 D0	01-01-1987
			ES 8702336 A1	16-03-1987
			ES 544402 D0	01-04-1988
			ES 8802131 A1	16-06-1988
			FI 834523 A ,B,	14-06-1984
			GB 2133788 A ,B	01-08-1984
			GB 2173787 A ,B	22-10-1986
			GR 79750 A1	31-10-1984
			IE 56324 B1	19-06-1991
			IL 70390 A	31-12-1986
			JP 1823303 C	10-02-1994
			JP 3178953 A	02-08-1991
			JP 5030826 B	11-05-1993
			JP 1762120 C	28-05-1993
			JP 3135948 A	10-06-1991
			JP 4040339 B	02-07-1992
			KR 9100436 B1	25-01-1991
			LU 88750 A9	23-08-1996
			MX 155545 A	25-03-1988
			PH 20074 A	18-09-1986
			PT 77771 A ,B	01-01-1984
			US 4761501 A	02-08-1988
			US 4611078 A	09-09-1986
			AU 2212383 A	21-06-1984